

Location/Identification

MINFILE Number:	092INW054	National Mineral Inventory Number:	092111 Gyp1
Name(s):	SPATSUM HART, FLORA, MARIE, BELLE, TOM, LOFAR, ORION, HIFAR, SOFAR, MARS		
Status:	Prospect	Mining Division:	Kamloops
Regions:	British Columbia	Electoral District:	Fraser-Nicola
BCGS Map:	0921054	Resource District:	Thompson Rivers Natural Resource District
NTS Map:	092111W	UTM Zone:	10 (NAD 83)
Latitude:	50 33 28 N	Northing:	5602025
Longitude:	121 18 11 W	Easting:	620199
Elevation:	425 metres		
Location Accuracy:	Within 500M		
Comments:	Adit, just west of Highway 1 and opposite the Spatsum station on the Canadian Pacific Railway, about 18.5 kilometres south of the community of Ashcroft (Assessment Report 6918).		

Mineral Occurrence

Commodities:	Copper, Zinc, Lead, Gypsum		
Minerals	Significant:	Chalcopyrite, Sphalerite, Galena, Gypsum	
	Alteration:	Gypsum, Pyrite, Silica, Barite, Talc	
	Alteration Type:	Silicific'n, Leaching, Pyrite	
	Mineralization Age:	Unknown	
Deposit	Character:	Stratiform, Disseminated, Massive	
	Classification:	Hydrothermal, Volcanogenic, Industrial Min.	
	Type:	G06: Noranda/Kuroko massive sulphide Cu-Pb-Zn	

Host Rock

Dominant Host Rock:	Metavolcanic		
Stratigraphic Age	Group	Formation	Igneous/Metamorphic/Other
Permian-Triassic	Undefined Group	Kutcho	-----
Isotopic Age	Dating Method	Material Dated	
-----	-----	-----	
Lithology:	Rhyolite Tuff, Rhyolite, Dacite Tuff, Dacite, Granite, Andesite Tuff, Andesite, Limestone, Diorite, Chert		

Geological Setting

Tectonic Belt:	Intermontane	Physiographic Area:	Thompson Plateau
Terrane:	Quesnel, Cache Creek		
Metamorphic Type:	Regional		
Grade:	Greenschist		

Inventory

No inventory data

Capsule Geology

Felsic volcanic and intrusive rocks which occur between the Martell and Bonaparte faults, near Ashcroft, are tentatively correlated with the Permo-Triassic Kutcho Assemblage, rather than the Upper Triassic to Lower Jurassic Nicola Group. Mafic volcanic rocks assigned to the Nicola Group occur both to the east and west of the Bonaparte fault. The presence of Upper Triassic fossils imply that this correlation is valid for basaltic rocks which occur east of the Bonaparte fault. However, the age of basaltic rocks that occur west of the Bonaparte fault, in proximity to, and possibly interbedded with rhyolite tuffs, is not constrained. These basaltic rocks may be contemporaneous with Lower Triassic felsic rocks, rather than the younger Nicola Group lavas. The presence of rocks of Kutcho Assemblage age and affinity raises the potential for Kutcho Creek-equivalent Cu-Zn volcanogenic massive sulphide mineralization (Fieldwork 1996).

The Spatsum property covers calcalkaline andesite to rhyolite metavolcanics and related chemical and clastic metasediments. They occur as a north to north-northwest striking, west dipping monoclinical sequence that has been metamorphosed to the mid-greenschist facies. The volcanics and sedimentary units have been locally intruded by diorite, granite, dacite and rhyolite plugs and dikes. Most of the intrusions are thought to be subvolcanic equivalents of the volcanic units. The metavolcanics include primarily andesite, dacite and rhyolite tuffs and tuff breccias and the metasediments consist of thin limestone and chert beds.

The most significant mineralization occurs where the rhyolite and locally the dacite units have been variably leached, silicified and pyritized and impregnated with gypsum, trace talc and barite, and very small amounts of chalcopyrite, sphalerite and galena. The mineralized and altered zones are heavily gossaned and pyrite concentrations are difficult to estimate due to the intense surface weathering and leaching. Gypsum occurs in significant concentrations in the zones as massive and/or disseminated clots commonly distributed throughout the altered rhyolite pyroclastics. Two mineralized gypsum-rich zones are about 600 metres apart and stand out prominently as large white masses. The larger and more southerly gypsum outcrop occurs over a strike length of 60 metres and a vertical height of 90 metres and strikes north-northeast with a moderate dip to the northwest. In 1913, an 8-metre exploratory adit was driven at the base of the southerly exposure; from the end of the adit a winze was sunk to a depth of 9 metres. The adit intersected a band of nearly pure white massive gypsum, 1.5 metres wide, which analysed 32.70 per cent CaO, 46.72 per cent SO₃, 20.60 per cent H₂O and 0.04 per cent insolubles (CANMET Report 714).

The alteration zones, because of the abundance of gypsum, are interpreted to represent a facies which commonly develops adjacent to many base metal-bearing Kuroko-type massive sulphide deposits. It is felt that base metal-bearing massive sulphide concentrations may exist along strike or downdip adjacent to these alteration zones. The only other mineralization observed on the property includes minor disseminated pyrite which occurs locally in some rhyolite units (Assessment Report 6918).

The Spatsum gypsum showings were first staked about in 1896 by a prospector named Munro, who did a small amount of development work, but allowed the lease to lapse. It was then taken up about 1906 by Messrs. Sinclair and Spencer, who staked four mineral claims called the Hart, Flora, Marie and Belle; the claims were surveyed in 1907. Work conducted by the El Paso Mining and Milling Company in 1972 comprised geological mapping and a soil geochemical survey. In 1978, D.H. Wilson prospected the Orion claim which covered the gypsum showings. In 1978-80, geological mapping, soil geochemical survey, induced polarization, VLF-EM and magnetic surveys, and percussion drilling of eight holes totalling 594 metres were performed by Cominco Ltd. In 1987-88, geological mapping and an induced polarization survey were completed on behalf of P.G. Dasler who interprets that the gypsum showings are vein gypsum and part of the zoning of an epithermal event. In 1995, prospecting was performed by D. Javorsky and F.M. Smith to evaluate the epithermal potential.

Bibliography

EMPR AR 1907-L134,L135; 1912-K185; 1922-N153; 1926-A194; 1962-42,43
EMPR ASS RPT 3680, *6918, 7102, 7638, 8263, 16963, 18388, 24280
EMPR EXPL 1978-E287,E288; 1979-332,333
EMPR FIELDWORK 1981, pp. 270,271; 1996, pp. 117-123
EMPR OF 1991-15; 1999-2
EMPR PF (Prospectors Report 1995-1 by Dave Javorsky, Prospectors Report 1996-8 by David Javorsky; Prospectors Report 1999-13 by David Javorsky)
EMPR PFD 811586, 823995
GSC MAP 1010A; 1386A; 42-1989
GSC MEM 262, p. 110
GSC OF 165; 866; 980
GSC P 46-8; 47-10; 69-23; 73-1A, p. 212; 74-49; 81-1A, pp. 185-189, 217-221; 82-1A, pp. 293-297; 85-1A, pp. 349-358
CANMET RPT *245 pp. 95-97; *714 pp. 63,64
CJES Vol.15, No.1 (January 1978), pp. 99-116
Grette, J.F. (1978): Cache Creek and Nicola Groups near Ashcroft, British Columbia, M.Sc. Thesis, University of British Columbia

Date Coded:	1985/07/24	Coded By:	BC Geological Survey (BCGS)	Field Check:	N
Date Revised:	2025/01/02	Revised By:	Kerri Shaw (KLS)	Field Check:	N

